

Stellar mass limits

- Hydrostatic equilibrium is the key element.
- Combine ideal classical gas law: $P_c = \frac{\rho_c}{\bar{m}} kT_c$.

- with expression for central pressure to get:

$$kT_c = \left(\frac{\pi}{36} \right)^{1/3} G \bar{m} M^{2/3} \rho_c^{1/3}.$$

- Applied to a contracting protostar, shows that T_c rises steadily as ρ_c increases.
- Temperature will continue to rise until either
 - Thermonuclear fusion starts to regulate T_c ; or
 - Electrons in core become degenerate
- Condition for stardom is that fusion sets in before electron degeneracy prevents further contraction.

AS 3003

Stellar Physics

Maximum core temperature

- Suppose star contracts until e^- are degenerate in core but ions form a classical gas. Then:

$$\begin{aligned} P_c &= K_{\text{NR}} n_e^{5/3} + n_i kT_c \\ &= K_{\text{NR}} \left(\frac{\rho}{m_{\text{H}}} \right)^{5/3} + \frac{\rho}{m_{\text{H}}} kT_c \text{ for a pure H plasma.} \end{aligned}$$

- Use hydrostatic equilibrium to eliminate P_c :

$$kT_c = \left(\frac{\pi}{36} \right)^{1/3} G m_{\text{H}} M^{2/3} \rho_c^{1/3} - K_{\text{NR}} \left(\frac{\rho_c}{m_{\text{H}}} \right)^{2/3}.$$

- Has form

$$\begin{aligned} kT_c &= A \rho_c^{1/3} - B \rho_c^{2/3} \text{ which has maximum value} \\ (kT_c)_{\text{max}} &= A^2/4B \text{ at } \rho_{c,\text{max}} = (A/2B)^3. \end{aligned}$$

AS 3003

Stellar Physics

Minimum mass of a star

- **Substituting, get**

$$(kT_c)_{\max} = \left(\frac{\pi}{36}\right)^{2/3} \frac{G^2 m_H^{8/3}}{4K_{\text{NR}}} M^{4/3}.$$

- **Need this max temp to be $> T_{\text{ign}}$, so:**

$$M_{\min} = \left(\frac{36}{\pi}\right)^{1/2} \left(\frac{4K_{\text{NR}}}{G^2 m_H^{8/3}}\right)^{3/4} (kT_{\text{ign}})^{3/4}.$$

$$\text{For } T_{\text{ign}} = 1.5 \times 10^6 \text{ K, } M_{\min} \approx 0.1 M_{\text{Sun}}.$$

AS 3003

Stellar Physics

Radiation pressure in stellar core

$$P_c = P_{\text{gas}} + P_{\text{rad}}, \text{ more conveniently written}$$

in terms of a parameter β :

$$P_{\text{gas}} = \beta P_c = \frac{\rho_c}{\bar{m}} kT_c \text{ and } P_{\text{rad}} = (1 - \beta)P_c = \frac{1}{3} a T_c^4.$$

Use β to eliminate T_c :

$$P_c = \left[\frac{3(1 - \beta)}{a \beta^4} \right]^{1/3} \left[\frac{k \rho_c}{\bar{m}} \right]^{4/3}. \text{ Equate this to}$$

hydrostatic equilibrium pressure :

$$\left[\frac{\pi}{36} \right]^{1/3} G M^{2/3} = \left[\frac{3(1 - \beta)}{a \beta^4} \right]^{1/3} \left[\frac{k \rho_c}{\bar{m}} \right]^{4/3}.$$

AS 3003

Stellar Physics

Maximum mass of a star

- Can use this equation to plot radiation pressure fraction $(1-\beta)$ as function of stellar mass.
- When $(1-\beta) > 0.5$ or so, radiation pressure destabilizes star -- so stars with $M > 50 M_{\text{Sun}}$ or so tend to blow apart

AS 3003

Stellar Physics

Understanding the main sequence

- Develop dimensionless equations of stellar structure for homologous models.
- Discover how mass-radius and mass-luminosity relations depend on opacity and energy generation.
- Determine slope of main sequence.
- Learn how changes in abundance affect location of main sequence in HR diagram.

AS 3003

Stellar Physics

Dimensionless structure equations

- **Define** $x = \frac{r}{R}$, $q = \frac{M(r)}{M}$, $t = \frac{T}{T_0}$, $p = \frac{P_g}{P_0}$.
- **and use** $\rho = \frac{\mu m}{k} \frac{P_g}{T}$.
- **Hydrostatic equilibrium becomes:**

$$\frac{P_0}{R} \frac{dp}{dx} = \frac{\mu m}{k} \frac{P_0}{T_0} \frac{p}{t} \frac{GM}{R^2} \frac{q}{x^2} \Rightarrow \frac{dp}{dx} = \frac{p}{t} \frac{q}{x^2}$$

$$\text{where } T_0 \equiv \frac{\mu m}{k} \frac{GM}{R}.$$

AS 3003

Stellar Physics

Mass continuity + energy transport

- **Similarly:** $\frac{M}{R} \frac{dq}{dx} = 4\pi R^2 x^2 \frac{\mu m}{k} \frac{P_0}{T_0} \frac{p}{t} \Rightarrow \frac{dq}{dx} = x^2 \frac{p}{t}$
 where $P_0 \equiv \frac{GM^2}{4\pi R^4}$.
- **Transport:** Let $\kappa = \kappa_0 \rho^\alpha T^{-\beta} = \kappa_0 \left(\frac{\mu m}{k} \frac{P_0 p}{T_0 t} \right)^\alpha (T_0 t)^{-\beta}$.

$$\frac{T_0}{R} \frac{dt}{dx} = - \frac{3\kappa_0}{16\pi ac (T_0 t)^3} \left(\frac{\mu m}{k} \frac{P_0 p}{T_0 t} \right)^{1+\alpha} (T_0 t)^{-\beta} \frac{L f}{R^2 x^2}$$

$$\Rightarrow \frac{dt}{dx} = -C \frac{p^{1+\alpha}}{t^{4+\alpha+\beta}} \frac{f}{x^2} \text{ where } C = \frac{3\kappa_0}{16\pi ac} \left(\frac{\mu m}{k} \right)^{1+\alpha} \frac{P_0^{1+\alpha}}{T_0^{5+\alpha+\beta}} \frac{L}{R}.$$

AS 3003

Stellar Physics

Energy generation

- **Energy generation:**

$$\text{Let } \varepsilon = \varepsilon_0 \rho T^\nu = \varepsilon_0 \frac{\mu m}{k} \frac{P_0 p}{T_0 t} (T_0 t)^\nu.$$

$$\frac{L}{R} \frac{df}{dx} = 4\pi r^2 \varepsilon \rho = 4\pi R^2 x^2 \varepsilon_0 \left(\frac{\mu m}{k} \frac{P_0 p}{T_0 t} \right)^2 (T_0 t)^\nu$$

$$\Rightarrow \frac{df}{dx} = D \frac{x^2 p^2}{t^{2-\nu}} \text{ where } D = \frac{4\pi \varepsilon_0 R^3}{L} \left(\frac{\mu m}{k} \right)^2 \frac{P_0^2}{T_0^{2-\nu}}.$$

AS 3003

Stellar Physics

Mass-radius relation: 1

- The constants P_0 , T_0 , C and D are unique to each star and apply throughout the entire structure.
- The product CD eliminates L to give a direct relation between mass and radius:

$$\begin{aligned} CD &= \frac{3\kappa_0 \varepsilon_0 R^2}{4ac} \left(\frac{\mu m}{k} \right)^{3+\alpha} \frac{P_0^{3+\alpha}}{T_0^{7+\alpha+\beta-\nu}} \\ &= \frac{3\kappa_0 \varepsilon_0 R^2}{4ac} \left(\frac{\mu m}{k} \right)^{3+\alpha} \cdot \left(\frac{GM^2}{4\pi R^4} \right)^{3+\alpha} \left(\frac{\mu m}{k} \frac{GM}{R} \right)^{\nu-7-\alpha-\beta} \\ &= \frac{3\kappa_0 \varepsilon_0}{4ac} \left(\frac{m}{k} \right)^{3+\alpha} \left(\frac{G}{4\pi} \right)^{3+\alpha} \left(\frac{mG}{k} \right)^{\nu-7-\alpha-\beta} \frac{\mu^{\nu-4-\beta} M^{\nu-1+\alpha-\beta}}{R^{\nu+3+3\alpha-\beta}} \end{aligned}$$

AS 3003

Stellar Physics

Mass-radius relation: 2

- Hence get mass-radius relation:

$$R^{\nu+3+3\alpha-\beta} \propto M^{\nu-1+\alpha-\beta}.$$

- Kramers' opacity law and CNO cycle give $\alpha=1$, $\beta=7/2$, $\nu=16$, so:

$$R \propto M^{12.5/18.5}.$$

- Böhm-Vitense finds that in practice, an opacity law with $\alpha=0.5$, $\beta=2.5$ gives a better fit:

$$R \propto M^{13/18}.$$