

Atomic processes :

Bound-bound transitions (Einstein coefficients)

Radiative processes from electron transitions:

- **Bound-bound:** electron moves between two bound states in an atom or ion. Photon emitted or absorbed.

$$h\nu = \chi_u - \chi_l$$

- **Bound-free:** electron moves between bound and unbound states. Bound-unbound: ionization. Unbound-bound: recombination

$$h\nu = \chi_{\text{ion}} - \chi_n + \frac{1}{2}mu^2$$

- **Free-free:** Free electron gains energy by absorbing a photon as it passes an ion, or loses energy by emitting a photon. This emission process is called Bremsstrahlung (braking).

$$h\nu = \frac{1}{2}mu_2^2 - \frac{1}{2}mu_1^2$$

Transition between two atomic energy levels:

Photon frequency, $h\nu_{ij} = |E_i - E_j|$

Hydrogenlike atoms (nucleus + one electron):

$$E_n = -Z^2 \frac{m_e e^4}{2n^2 \hbar^2} \equiv -\frac{Z^2 R}{n^2}$$


where

n is an integer (the principal quantum number),
 Z is nuclear charge in units of e , and

$R \cong 13.6 \text{ eV}$ is a constant.

Spectrum consists of a series of lines, labelled by the final n of downward transition. eg. the Lyman series are transitions to $n=1$.

Lyman α is the transition $n=2$ to $n=1$,
wavelength $\lambda(\text{Ly}\alpha) = 121.6 \text{ nm}$.



Boltzmann's Law

- In thermodynamic equilibrium at temperature T , the populations n_1 and n_2 of any two energy levels are given by Boltzmann's law,

$$\frac{n_2}{n_1} = \frac{g_2}{g_1} e^{-(E_2 - E_1)/kT}$$

- E_1 and E_2 are the energies of the levels relative to the ground state.
- Some energy levels are degenerate (i.e. can hold >1 electron). Statistical weights g_1, g_2 give the number of sublevels.
- In terms of photon frequency:

$$\frac{n_2}{n_1} = \frac{g_2}{g_1} e^{-h\nu/kT}$$

HYDROGEN ATOM

Excitation energy

$$\chi_n = \chi_{ion} \left(1 - \frac{1}{n^2}\right)$$

Statistical weight of level n is $2n^2$

$n=1$, Lyman series 1216- 912 Å


$n=2$, Balmer series 6563-3647 Å

$n=3$, Paschen series 18751-8204 Å

$n=4$, Brackett series 40512-14584 Å

$n=5$, Pfund series 74578-22788 Å

Astrophysical Formulae, Lang



Bound-bound transitions: Einstein coefficients

- Kirchhoff's Law relates the absorption and emission coefficients for black body radiation,


$$B_{\nu} = \frac{j_{\nu}}{\alpha_{\nu}}$$

- This law
 - was derived without using any knowledge of microscopic processes.
 - Must imply some relation between emission and absorption processes at an atomic level.

2-level atom

- Einstein considered the case of a two level atom:

- Two energy levels,
- Energy E_1 , statistical weight g_1 .
- Energy $E_1 + \Delta E = E_1 + h\nu_0$, statistical weight g_2 .
- 3 important radiative processes follow.



1. Spontaneous emission

- Atom decays spontaneously from level 2 to level 1.
- Photon emitted.
- Occurs independently of the radiation field.
- **Define:** The Einstein A -coefficient, A_{21} , is the transition rate per unit time for spontaneous emission ($\sim 10^8 \text{ s}^{-1}$).

2. Absorption


- Photons with energies close to $h\nu_0$ cause transitions from level 1 to level 2.
- The probability per unit time for this process will evidently be proportional to the mean intensity at the frequency ν_0 .

Line profile $\phi(\nu)$

Need to define a *line profile function*

$\phi(\nu)$:

- describes the probability that a photon of frequency ν will cause a transition.
- $\phi(\nu)$ is sharply peaked at ν_0 , with width $\Delta\nu$ and normalization, $\int_0^\infty \phi(\nu) d\nu = 1$



Define: The transition rate per unit time for absorption is $B_{12}\bar{J}$

where, $\bar{J} \equiv \int_0^\infty J_\nu \phi(\nu) d\nu$

with J_ν being the mean intensity and $\phi(\nu)$ the line profile function.

B_{12} is one of the Einstein B-coefficients.

Note: we have been careful to distinguish between J_ν and \bar{J} , but this is a technicality. If J_ν changes slowly over the line width $\Delta\nu$ of the line, then $\phi(\nu)$ is almost $\delta(\nu - \nu_0)$ and $\bar{J} \cong J_{\nu_0}$

3. Stimulated emission

Planck's law does not follow from considering only spontaneous emission and absorption. Must also include *stimulated emission*, which like absorption is proportional to \bar{J}

Define: $B_{21}\bar{J}$ is the transition rate per unit time for stimulated emission.

B_{21} is a second Einstein B-coefficient. Stimulated emission occurs into the same state (frequency, direction, polarization) as the photon that stimulated the emission.

Lecture 6 revision quiz

- Calculate the wavelengths of the first 3 lines of the hydrogen Balmer series: $H\alpha$, $H\beta$, $H\gamma$.
- Define the statistical weight g of an atomic energy level.
- Write down Boltzmann's Law and define all symbols used and their units.